



e la Facultad de Medicina Veterinaria y Zootecnia http://veterinariamexico.unam.mx/

Efficacy of *Metarhizium anisopliae* in the control of the horn fly, *Haematobia irritans* (*Diptera: Muscidae*), under natural infestation conditions

Abstract

The aim of this study was to determine the efficacy of the entomopathogenic fungus, *Metarhizium anisopliae* strain 134 (Ma134), for dairy cattle naturally infested with *Haematobia irritans* (L.) An aqueous formulation of 1×10^8 conidia/ml was applied by aspersion on a group of eight cows every seven days for a total of four applications, while a control group was sprayed with the same solution without conidial content. The degree of horn fly infestation was estimated daily in both groups by the direct count of adult flies found resting or feeding on the animals, and Abbott's formula was used to calculate the effectiveness of the formulation. The Ma134 formulation resulted in a 68.6 % reduction of the fly infestation during the study period, and a significant reduction was observed from the first week post-application (p < 0.05). The results show that the Ma134 formulation is effective in reducing infestations of horn fly, *H. irritans*, on dairy cattle under natural conditions. Thus, the fungus, Ma134, can be considered a candidate for biological control for the integrated management of *H. irritans*.

Keywords: Haematobia irritans, Metarhizium anisopliae, natural infestation, biological control.

Introduction

The horn fly, *Haematobia irritans* (L.), is common and primarily affects cattle on grass pastures in Mexico. The horn fly was introduced from Europe to North America in the late 19th century. Currently, it is widely distributed across the American continent and is considered the major pest of livestock.^{1,2} *H. irritans* is an obligate bloodsucking ectoparasite, always remains on cattle and feeds 24 to 38 times a

Cite this as:

Cruz-Vázquez C, Carvajal-Márquez J, Lezama-Gutiérrez R, Vitela-Mendoza I, Ángel-Sahagún CA. Efficacy of *Metarhizium anisopliae* in the control of the horn fly, *Haematobia irritans (Diptera: Muscidae)*, under natural infestation conditions. Veterinaria México OA. 2017;4(2). doi: 10.21753/vmoa.4.2.384.

Carlos Cruz-Vázquez^{1*} 0000-0003-0970-6811 Juan Carvajal Márquez¹ 0000-0002-9831-2381 Roberto Lezama Gutiérrez² 0000-0002-5192-2895 Irene Vitela-Mendoza¹ 0000-0002-7592-6102 César Andrés Ángel-Sahagún³ 0000-0001-5370-6400

¹ División de Estudios de Posgrado e Investigación. Instituto Tecnológico El Llano Aguascalientes, km 18 carretera Aguascalientes- San Luis Potosí, 20330, El Llano, Aguascalientes, México.

² Facultad de Ciencias Biológicas y Agropecuarias. Universidad de Colima, km 40 carretera Colima-Manzanillo, 28100, Tecomán, Colima, México.

³ División de Ciencias de la Vida, Campus Irapuato-Salamanca, Universidad de Guanajuato, exhacienda El Copal, km 7, Carretera Irapuato-Silao, 36500, Irapuato, Guanajuato, México.

> *Corresponding author: Tel: +52-449-916-12-51 Email address: cruva18@yahoo.com.mx

Received: 2016-05-12 Accepted: 2017-03-31 Published: 2017-04-06 Additional information and declarations

can be found on page 7

© Copyright 2017 Carlos Cruz-Vázquez et al.





Distributed under Creative Commons CC-BY 4.0

day, consuming an average of 1.71 mg of blood per meal.² The feeding habits of horn flies on cattle cause irritation and discomfort. In an effort to repel the flies, the animals develop defensive behaviours such as tail flicks, leg stomps, skin twitches and head throws. This stress, which varies in intensity based on the quantity of flies, results in reduced grazing time and increased energy spent.³⁻⁵ Hence, treating horn fly infestations are of sanitary and economic importance.

It has been estimated that weight loss can be up to 220 g/day and milk production can be between 10 and 20 % of regular production when the infestation is high (> 300 flies/animal). Furthermore, *H. irritans* is an efficient transmitter of several viral, bacterial, protozoal, and rickettsial diseases and helminth infections.^{2,6,7} The bites of the horn fly reduce the quality of the hides, and insecticide application is costly. In general, control measures of the horn fly for grazing cattle include applying insecticides of different chemicals through the use of earrings, aspersions, immersion baths and pour-on formulations; however, the resistance to insecticides is present in several livestock regions of South America.⁸⁻¹⁰ In Mexico, the problem is more severe in tropical regions.¹¹⁻¹⁴

The use of insecticides has a considerable impact on public health due to the contamination of meat and milk with toxic residues and through environmental pollution. An alternative for reducing the intensive use of insecticides is to use non-chemical measures such as biological control. For example, the entomopathogenic fungi is a safe and sustainable alternative with minimum risk for vertebrates, humans and the environment. This fungus infects the insect through contact, invading it through the cuticle and killing it. After the death of the insect, the fungus emerges from the cadaver to produce new conidia.¹⁵

Currently, the efficacy of the fungus *Metarhizium anisopliae* as a biological control of *H. irritans* has been demonstrated under laboratory conditions in both immature and adult stages,¹⁶⁻¹⁹ as well as in adults under controlled infestation conditions.²⁰ However, the efficiency of this entomopathogenic fungus has been scarcely studied under field conditions.²¹ To advance the development of this biological control, the objective of this study was to determine the effectiveness in reducing the infestation of *H irritans* on dairy cattle under natural conditions, by applying the strain Ma134 of *M. anisopliae* by aspersion.

Material and methods Study site

This work was performed at the livestock unit of El Llano Technological Institute, located in a municipality of the same name in Aguascalientes, Mexico. The place is 2 020 masl, with a semiarid climate, summer rains, and an average temperature of 15.5 °C. In the region, natural infestation of *H. irritans* is seasonal, being the highest from summer to autumn.^{22,23}

Entomopathogenic fungi

The strain *M. anisopliae sensu lato* 134 (Ma134) was isolated from soil in the dairy unit at El Llano Technological Institute using the larvae of *Galleria mellonella* L. (*Lepidoptera: Pyralidae*) as a reservoir host. Taxonomic identification was based



on morphological criteria of the reproductive structures; the strain forms part of the entomopathogenic fungi collection at the same Institution.²⁴ The strain Ma134 was evaluated in *Musca domestica* and *Stomoxys calcitrans* adults under *in vitro* conditions with good results²⁴ and high efficacy with *S. calcitrans* under field conditions.²⁵ The strain was cultured in the laboratory according to the protocol previously described by Cruz-Vázquez et al.²⁵ The isolate was cultured on Sabouraud's Dextrose Agar enriched with 1 % yeast extract²⁶ containing 500 ppm chloramphenicol and incubated at 25 ± 1 °C for 21 days in a 12:12 hr light/dark regime. Conidia were harvested by scraping, suspended in sterile distilled water containing 0.1 % (v:v) Tween 80, and homogenized on a vortex mixer. Spore viability, which exceeded 98 %, was determined by seeding 100 µl of conidial suspension on Sabouraud's Dextrose Agar and counting colonies 48 h later. Mass reproduction was performed on grains of rice according to that described by Ángel-Sahagún et al.²⁷

Field trial

Two experimental groups were formed with eight Holstein cows, each group was maintained in an area of 5 000 m2, separated from each other by 500 m. Both areas had ryegrass pasture (Lolium perenne), water tanks for drinking and natural shade provided by various native trees. The cows did not receive any treatment to control fly infestation during the year prior to this trial. The experiment began the first week of September, before the second fly population peak of the season according to information reported in the area.^{22,23} The first group received an application of the aqueous formulation of Ma134 fungi (1 x 10⁸ conidia/ml concentration), water, Polyoxyethylene sorbitan monooleate (Tween 80®, Sigma-Aldrich Co., St. Louis, MO, USA) solution (0.01 %) and an agricultural adjuvant at 0.1 % (Inex-A, Cosmocel, Mexico). Five litres of the formulation were applied per animal by aspersion using a back sprayer with a cone-type nozzle at a pressure of approximately 40 lb/in2, on four occasions with seven-day intervals between treatments. The formulation was prepared 30 min prior to administration. The control group received a water solution with Tween 80 (0.01 %) plus an agricultural adjuvant at 0.1 %. Sprayings were between 7 and 8 pm to avoid sunlight. The number and interval of applications were similar to that used in a previous study with the stable fly.²⁵ The Committee on Use and Care of Animals of the Instituto Tecnológico El Llano Aguascalientes approved this project, and adequate veterinary care was provided to all of the animals under study.

Estimation of H. irritans infestation

An Infestation index (average number of flies per animal) was estimated daily by the direct count of adult flies found resting or feeding on the animals; each cow was considered an experimental unit. This estimation consisted of photographing only one side of the body of each animal (including regions of the head, neck, back, sides and limbs). using a digital camera with a 5x optical zoom function. This activity was done daily between 8 and 9 am, always by the same person, who was not aware of the status of the cows. Counting the number of flies was performed by computer image analysis. The obtained number was multiplied by two to obtain



Original Research



Figure 1. Infestation index of horn flies, Haematobia irritans, on cows treated with the aqueous formulation of Ma134

an estimate of the total number of flies.^{20,28} Climatic factors were registered as reported by the El Llano weather station, the measurements included the average daily temperature (T °C), the average daily relative humidity (RH %) and the rainfall (Rf) during the study period.

Data Analysis

The information recorded in the groups under study (treated and control) was analysed by analysis of variance and Student's t-test (p < 0.05), with the objective to demonstrate the differences between the groups in each study week using the corresponding week's average index of infestation values. Then, Abbott's formula was used to determine the efficacy of the formulation²⁹ by identifying the percent reduction in the average infestation index for each evaluated week.²⁰

Results and Discussion

The population curve of the horn fly, *H. irritans*, in the groups under study is shown in figure 1. On the second day post-treatment, the daily index infestation began to decrease compared to the control group, maintaining this trend until the end of the study. Infestation index at day 0 in the treated group was 530 and in the control group, 532. The aqueous formulation reduced the index infestation of 538, which was determined at the start of work (day 1), to 169 on day 28 post-treatment, representing 68.6 % efficiency.

In Table 1, the average infestation index is shown in each of the study weeks. The infestation index was reduced from week to week in the treated group. In the **Table 1.** Infestation index average and efficacy of the Ma134 aqueous formulation applied on Holstein cows naturally
infested with *H. irritans*. Different subscripts (a,b) on the same row indicate significant differences (p < 0.05).
Efficacy: reduction percentage of the average infestation index calculated with Abbott's formula.

Study week	Treated group	Control group	Efficacy (%)
1	465 ± 20.8a	530 ± 25.2b	12.26
2	378 ± 21.1a	562 ± 17.2b	32.60
3	296 ± 9.6a	569 ± 14.7b	47.98
4	221 ±14.8 a	479 ± 10.7b	53.86

control group, it was maintained between 530 and 569 in the first three weeks; however, in the fourth week, it was reduced to 479. The number of flies in the treated group was always less than in the control group. Statistical differences were detected each week between treated and control groups: first week ($t_{14} = -3.14$, p = 0.006), second week ($t_{14} = -10^{-19}$, $p = 7.3 \times 10^{-8}$), third week ($t_{14} = -25.88$, $p = 3.17 \times 10^{-13}$), and fourth week ($t_{14} = -9.08$, $p = 3.04 \times 10^{-7}$). The cattle remained healthy during the study period. Climatic data during the study period indicated that the average temperature was 20.2 °C with a range from 18 to 21.6 °C, while the RH was on average 59.7 % with a range from 38.4 to 80.3 %, and no rainfall was registered (figure 2).

The results obtained in this study show that the formulation of the Ma134 strain was able to control horn fly infestation under field conditions. Compared to the control group, an infestation reduction of the fly in the treatment group was consistent during the study, and the weekly applications helped reinforce the pathogenic action of the fungus. Thus, it can be considered that the treatment had an additive effect. The aqueous formulation of Ma134 was added to an agricultural adjuvant to promote adhesion of conidia to the flies and animal hair and provide assistance in protecting the conidia from exposure to sunlight. In the literature, it is mentioned that conidia are sensitive to UV rays and to high environmental temperatures, and when applied to the body of animals conidia are sensitive to body temperature.³⁰ The design of a formulation for topical application should consider at a minimum these factors; hence, the incorporation of an agricultural adjuvant or another compound with a heat/UV protecting ability can improve the performance of the conidia.^{21,27} However, Angel-Sahagún et al.²⁷ did not find that environmental conditions had any effect on the performance of entomopathogenic fungi used in experimental grasslands artificially infested with larvae of *Rhipicephalus microplus*. The tolerance of entomopathogenic fungi to environmental conditions appears to be related to the site where the fungus was isolated,³¹ and it can be assumed that the Ma134 strain is adapted to a semiarid climate, the same as the study site, which probably contributed to its good performance in the study.

M. anisopliae has been the most studied entomopathogenic fungus because of its cosmopolitan and highly pathogenic nature. In addition to the fungus being specifically pathogenic for common insects affecting domesticated animals, such as ticks and flies, it is an organism that has minimum risk for plants, animals, humans and the environment.^{15,30,32} In this study, no adverse effects of the fungus were observed in the cattle. Other authors reported similar findings in cattle infested with ticks or stable flies and treated with *M. anisopliae* formulations under field conditions.^{25,33,34} The homogenous weather conditions in the study period and an ad-



Figure 2. Climatic conditions in the study period: Relative Humidity (RH) and Temperature (°C)

equate combination of humidity and temperature were ideal for the development and maintenance of the population curve of *H. irritans*. However, at the end of the study, the change in these environmental conditions, especially the low average daily temperature, favoured a decline in the infestation of flies. Cruz-Vázquez et al.^{22,23} describe similar population dynamics in the region, characterized by a close relationship with temperature and relative humidity. In these studies, it is mainly suggested that temperature induces a decline in population growth and in the late fall temperature can cause diapause.

Recently, in Tecoman, Colima, Mexico, a region with a tropical dry climate, the effect of several strains of *M. anisopliae* on steers with an experimental infestation of *H. irritans* was reported: mortality was observed from day 1 post-treatment (20 %), on day 4 it was 61 %, by day 8 it was 84 %, and from days 12 to 13, it was between 94 % and 100 %. However, the absence of reinfestation limited the possibilities of estimating the longevity of the effect of the fungus.²⁰ In São Paulo state, Brazil, a region with a sub-tropical climate, a study was performed to evaluate the susceptibility of adult *H. irritans* flies to a water-based formulation of a strain of *M. anisopliae* under natural infestation conditions. For this purpose, the formulation had a concentration of 3 x 10¹⁰ conidia /ml and was applied to cattle every five days on four occasions. This study found that the level of infestation reduced in the treated group from the second spraying of the formulation, while in the untreated group infestation always increased. At the end of the study, the treated group had an average of 28.5 flies/head while the untreated group had 65.4 flies/head.²¹

In our study, an additive effect was also found. This phenomenon was observed both in cattle naturally infested with *S. calcitrans*²⁵ and in cattle infested with *R. microplus* ticks under field conditions.³⁴ The use of multiple applications of an entomopathogenic fungus provides better results.^{30,33} The results obtained in this study confirm the potential of *M. anisopliae* to infect and cause death in *H. irritans* under natural infestation conditions. Because the infestation by the horn fly in the region under study is seasonal, it would be advisable to start a control programme using this formulation when the first detection occurs to limit the fly population growth. Thereby, reducing its adverse effects on production and animal welfare, as well as the use of insecticides. This biological control is a safe and sustainable alternative that has minimum risk for vertebrates, humans and the environment.^{15,30} According to the literature, it is possible to use in combination with other biological control agents, such as parasitoids³⁵ or insecticides.³⁶ To our knowledge, there are no studies on the simultaneous use of different natural enemies or combinations with insecticides in the case of the horn fly. Nevertheless, to transfer this alternative of biological control to ranchers, more research is necessary including (1) a completion of more field trials with a greater number of animals in different conditions to further validate the biological control's performance, and (2) an estimation of the cost-benefit ratio.

Conclusions

The results of this study demonstrated that the Ma134 formulation was 68 % effective in reducing infestation of the horn fly, *H. irritans*, on dairy cattle under natural infestation conditions. For this reason, the formulation can be considered a biological control for the integrated management of this insect. However, further investigation is needed to ensure its performance as a biological control.

Funding

This project was financed by ITEL-TecNM.

Acknowledgements

We gratefully acknowledge Dr. Miguel Ramos Parra (ITEL-TecNM) for his careful review of the statistical analysis of the study.

Conflicts of interest

None of the authors had any conflict of interest in regard to this article.

Author contributions

Carlos Cruz-Vázquez: conceived and designed the research, drafted and wrote the article.

Juan Carvajal Márquez: completed field and laboratory work, wrote the article.

Roberto Lezama Gutiérrez: developed the experimental design, supervised the experiment.

Irene Vitela-Mendoza: completed field and laboratory work.

Cesar Ángel-Sahagún: analysed the data.



References

- 1. Byford RL, Craig ME, Crosby BL. A review of ectoparasites and their effect on cattle production. J Anim Sci. 1992;70(2):597-602. doi: 10.2527/1992.702597x.
- Foil LD, Hogsette JA. Biology and control of tabanids, stable flies and horn flies. Rev - Off Int Epizoot. 1994;13:1125-58.
- 3. Harvey TL, Launchbaugh JL. Effect of horn flies on behavior of cattle. J Econ Entomol. 1982;75(1):25-7. doi: 10.1093/jee/75.1.25.
- 4. Harvey TL, Brethour JR. Effect of horn flies on weight gains of beef cattle. J Econ Entomol. 1979;72(4):516-8. doi: 10.1093/jee/72.4.516.
- Boland HT, Scaglia G, Umemura K. Case study impact of horn flies, *Haema-tobia irritans* (L.) (*Diptera: Muscidae*), on the behavior of beef steers. PAS. 2008;24(6):656-60. doi: 10.15232/S1080-7446(15)30918-9.
- Guglielmone AA, Gimeno E, Ediat J, Fisher WF, Volpogni MM, Quiano O, et al. Skin lesions and cattle hide damage from *Heamatobia irritans* infestations. Med Vet Entomol. 1999;13(3):324-9. doi: 10.1046/j.1365-2915.1999.00167.x.
- Cruz-Vázquez C. Infestación por moscas. In: Ibarra VF, Figueroa CJA, Quintero M, editors. Parasitología veterinaria volúmen III artrópodos. CDMX (MX): Universidad Nacional Autónoma de México; 2012. p. 61-70.
- Oyarzún MP, Quiroz A, Birkett MA. Insecticide resistance in the horn fly: alternative control strategies. Med Vet Entomol. 2008;22(3):188-202. doi: 10.1111/j.1365-2915.2008.00733.x.
- Guglielmone AA, Castelli ME, Volpogni MM, Medus PD, Martins JR, Suárez VH, et al. Toxicity of cypermethrin and diazinon to *Haematobia irritans* (*Diptera: Muscidae*) in its american southern range. Vet Parasitol. 2001;101(1):67-73. doi: 10.1016/S0304-4017(01)00490-3.
- Barros ATM, Alison JMW, Foil LD. Evaluation of a yearly insecticidal ear tag rotation for control of pyrethroid-resistant horn flies (*Diptera: Muscidae*). Vet Parasitol. 1999;82(4):317-25. doi: 10.1016/S0304-4017(99)00024-2.
- Taboada-Romero YL, Olivares-Pérez J, Gutiérrez-Segura I. Diagnosis of resistance *Haematobia irritans (Diptera: Muscidae*) to cypermethrin and coumaphos in bovines ranches of Tierra Caliente, Guerrero, Mexico. Revista Científica. 2013;23(4):283-6.
- Martínez IF, Osorio MJ, Ortiz NA, Fragoso SH, Giles HI. Current situation of pyrethoid resistance in the horn fly *Haematobia irritans* in ten mexican states. V international seminary in animal parasitology; Mérida, Yucatán, México 2003. p. 231-9.
- Kunz SE, Ortiz-Estrada M, Fragoso SH. Status of *Haematobia irritans* (*Diptera: Muscidae*) insecticide resistance in northeastern Mexico. J Med Entomol. 1995;32(5):726-9. doi: 10.1093/jmedent/32.5.726.
- Almazán GC, Cantú CA, Vega FA, García-Vázquez Z, Kunz S, Medellín LA. Situación de la resistencia a la cipermetrina y diazinon en mosca del cuerno (*Haematobia irritans*) en Tamaulipas, México. Veterinaria (Méx.) 2004;35(3):237-44. doi: 10.1093/jmedent/32.5.726.
- Zimmerman G. Review on safety of entomopathogenic fungus *Me-tarhizium anisopliae*. Biocontrol Sci Tech. 2007;17(9):879-920. doi: 10.1080/09583150701593963.



- Mochi DA, Monteiro AC, Ribeiro MAC, Yoshida L. Entomopathogenic fungal activity against pupae and adult *Haematobia irritans (Diptera: Muscidae)*. Vet Parasitol. 2010;168(1-2):105-10. doi: 10.1016/j.vetpar.2009.10.002.
- Mochi DA, Monteiro AC, Ribeiro MAC, Yoshida L. Efficiency of entomopathogenic fungi in the control of eggs and larvae of the horn fly *Haematobia irritans* (*Diptera: Muscidae*). Vet Parasitol. 2010;167(1):62-6. doi: 10.1016/j. vetpar.2009.09.046.
- Lohmeyer K, Miller J. Pathogenicity of three formulations of entomopathogenic fungi for control of adult *Haematobia irritans* (*Diptera: Muscidae*). J Econ Entomol. 2006;99(6):1943-7. doi: 10.1093/jee/99.6.1943.
- Ángel-Sahagún CA, Lezama-Gutiérrez R, Molina-Ochoa J, Galindo-Velasco E, López-Edwards M, Rebolledo-Domínguez O, et al. Susceptibility of biological stages of the horn fly, *Haematobia irritans*, to entomopathogenic fungi (*Hypho-mycetes*). J Insect Sci. 2005;5(1):50. doi: 10.1093/jis/5.1.50.
- Galindo-Velasco E, Lezama-Gutiérrez R, Cruz-Vázquez C, Pesador-Rubio A, Ángel-Sahagún CA, Ojeda-Chi MM, et al. Efficacy of entomopathogenic fungi (Ascomycetes:Hypocreales) against adult *Haematobia irritans* (*Diptera:Muscidae*) under stable conditions in the Mexican dry tropics. Vet Parasitol. 2015;209 (3-4):173-8. doi: 10.1016/j.vetpar.2015.02.025.
- Mochi DA, Monteiro AC, Simi LD, Moraes SA. Susceptibility of adult and larvae stages of the horn fly *Haematobia irritans*, to entomopathogenic fungus *Metarhizium anisopliae* under field conditions. Vet Parasitol. 2009;166(1-2):136-43. doi: 10.1016/j.vetpar.2009.07.037.
- Cruz-Vázquez C, Ramos PM, García-Vázquez Z, Perales SC, Mata D. Influencia de algunos factores climáticos en la distribución anual de la infestación por *Haematobia irritans (Diptera: Muscidae)* en un establo lechero de Aguascalientes, México. Veterinaria (Méx.) 2003;34(4):315-22.
- Cruz-Vázquez C, Bautista HJ, Vitela MI, Ramos PM, García-Vázquez Z, Quintero-Martínez T. Distribución anual de *Haematobia irritans* en tres establos lecheros en Aguascalientes, México. Veterinaria (Méx.) 2000;31(3):195-9.
- López-Sánchez J, Cruz-Vázquez C, Lezama-Gutiérrez R, Ramos-Parra M. Effect of entomopathogenic fungi upon adults of *Stomoxys calcitrans* and *Musca domestica* (*Diptera: Muscidae*). Biocontrol Sci Technol. 2012;22(8):969-73. doi: 10.1080/09583157.2012.699026.
- Cruz-Vázquez C, Carvajal-Márquez J, Lezama-Gutiérrez R, Vitela-Mendoza I, Ramos-Parra M. Efficacy of the entomopathogenic fungi *Metarhizium anisopliae* in the control of infestation by stable flies *Stomoxys calcitrans* (L.), under natural infestation conditions. Vet Parasitol. 2015;212(3-4):350-5. doi: 10.1016/j.vetpar.2015.07.003.
- Watson DW, Geden CJ, Long SJ, Rutz DA. Efficacy of Beauveria bassiana for controlling the house fly and stable fly (*Diptera: Muscidae*). Biol Control. 1995;5(3):405-11. doi: 10.1006/bcon.1995.1048.
- Ángel-Sahagún C, Lezama-Gutiérrez R, Molina-Ochoa J, Pescador-Rubio A, Skoda SR, Cruz-Vázquez C, et al. Virulence of mexican isolates of entomopathogenic fungi (*Hypocreales: Clavicipitaceae*) upon *Rhipicephalus=Boophilus microplus* (*Acari: Ixodidae*) larvae and the efficacy of conidia formulations under field conditions. Vet Parasitol. 2010;170(3-4):278-86. doi: 10.1016/j. vetpar.2010.02.037.



- Galindo-Velasco E, Cruz-Vázquez C, Lezama-Gutiérrez R, Reyes-Velázquez WP, Aguilar-Espinoza S, Pescador-Rubio A. Fluctuación poblacional de *Haematobia irritans* (*Diptera: Muscidae*) en un hato bovino en Tecomán, Colima, México. Veterinaria (Méx.) 2008;39(2):181–6.
- 29. Hillerton JE, Bramley AJ, Broom DM. The distribution of five species of flies (*Diptera: Muscidae*) over the bodies of dairy heifers in England. Bull Entomol Res. 1984;74(1):113-9. doi: 10.1017/S0007485300009986.
- Fernandes EKK, Bittencourt VR, Roberts DW. Perspectives on the potential of entomopathogenic fungi in biological control of ticks. Exp Parasitol. 2012;130(3):300-5. doi: 10.1016/j.exppara.2011.11.004.
- Rangel DEN, Braga GUL, Anderson AJ, Roberts DW. Variability in conidial thermotolerance of *Metarhizium anisopliae* isolates from different geographic origins. J Invertebr Pathol. 2005;88(2):116–25. doi: 10.1016/j.jip.2004.11.007.
- Ojeda-Chi MM, Rodriguez-Vivas RI, Galindo-Velasco E, Lezama-Gutiérrez R, Cruz-Vázquez C. Control de *Rhipicephalus microplus (Acari: Ixodidae)* mediante el uso del hongo entomopatógeno *Metarhizium anisopliae (Hypocreales: Clavivipitaceae)*. Revisión. Rev Mex Cienc Pecu. 2011;2(2):177-92.
- 33. Kaaya GP, Samish M, Hedimbi M, Gindin G, Glazeer I. Control of tick population by spraying *Metarhizium anisopliae* conidia on cattle under field conditions. Exp Appl Acarol. 2011;55(3):273-81. doi: 10.1007/s10493-011-9471-3.
- Alonso-Diaz MA, García L, Galindo-Velasco E, Lezama-Gutiérrez R, Angel-Sahagún CA, Rodriguez-Vivas RI, et al. Evaluation of *Metarhizium anisopliae* (*Hyphomycetes*) for the control of *Boophilus microplus* (*Acari: Ixodidae*) on naturally infested cattle in the Mexican tropics. Vet Parasitol. 2007;147(3-4):336–40. doi: 10.1016/j.vetpar.2007.03.030.
- Mesquita ALM, Lacey LA. Interactions among the entomopathogenic fungus, *Paecilomyces fumosoroseus (Deuteromycotina: Hyphomycetes)*, the parasitois *Aphelinus asychis (Hymenoptera: Aphelinidae)*, and their aphid host. Biol Control. 2001;22(1):51-9. doi: 10.1006/bcon.2001.0950.
- Bahiense TC, Fernandes EK, Bittencourt VREP. Compatibility of the fungus *Metarhizium anisopliae* and deltamethrin to control a resistant strain of Boophilus microplus tick. Vet Parasitol. 2006;141(3-4):319-24. doi: 10.1016/j. vetpar.2006.05.011.